


Ramachandran Plot

pdb3p4q




Plot statistics

Residues in most favoured regions [A,B,L]	1658	90.7%
Residues in additional allowed regions [a,b,l,p]	159	8.7%
Residues in generously allowed regions [-~a,-~b,-~l,-~p]	8	0.4%
Residues in disallowed regions	3	0.2%
-----		-----
Number of non-glycine and non-proline residues	1828	100.0%
Number of end-residues (excl. Gly and Pro)	16	
Number of glycine residues (shown as triangles)	188	
Number of proline residues	106	
-----		-----
Total number of residues	2138	

Based on an analysis of 118 structures of resolution of at least 2.0 Angstroms and R-factor no greater than 20%, a good quality model would be expected to have over 90% in the most favoured regions.

Ramachandran plots for all residue types


pdb3p4q



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Ramachandran plots for all residue types


pdb3p4q



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Ramachandran plots for all residue types


pdb3p4q



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Chi1-Chi2 plots


pdb3p4q



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Chi1-Chi2 plots


pdb3p4q



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Main-chain parameters

pdb3p4q




Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. %-tage residues in A, B, L	1828	90.7	57.8	10.0	3.3	BETTER
b. Omega angle st dev	2127	4.9	6.0	3.0	-0.4	Inside
c. Bad contacts / 100 residues	17	0.8	28.5	10.0	-2.8	BETTER
d. Zeta angle st dev	1950	1.5	3.1	1.6	-1.0	Inside
e. H-bond energy st dev	1376	0.9	1.1	0.2	-1.3	BETTER
f. Overall G-factor	2138	0.1	-0.9	0.3	3.4	BETTER

Side-chain parameters

pdb3p4q




pdb3p4q

Plot statistics


Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. Chi-1 gauche minus st dev	221	14.4	30.4	6.5	-2.5	BETTER
b. Chi-1 trans st dev	581	16.3	29.0	5.3	-2.4	BETTER
c. Chi-1 gauche plus st dev	838	15.5	27.7	4.9	-2.5	BETTER
d. Chi-1 pooled st dev	1640	15.7	28.5	4.8	-2.6	BETTER
e. Chi-2 trans st dev	488	18.1	27.7	5.0	-1.9	BETTER

Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




Residue properties

pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility



Key:- Helix Beta strand Random coil Accessibility shading: Buried Accessible


e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb3p4q


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors



Main-chain bond lengths

pdb3p4q




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

Main-chain bond angles

pdb3p4q




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

Main-chain bond angles

pdb3p4q




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

RMS distances from planarity

pdb3p4q




Histograms showing RMS distances of planar atoms from best-fit plane.
 Black bars indicate large deviations from planarity: RMS dist > 0.03 for rings, and > 0.02 otherwise.

Distorted geometry


pdb3p4q

Main-chain bond lengths



Bonds differing by > 0.05 Å from small-molecule values. Values shown: "ideal", difference, actual

Main-chain bond angles



Bond angles differing by > 10.0 degrees from small-molec values. Values shown: "ideal", actual, diff.