


Ramachandran Plot

pdb1nek



Plot statistics

Residues in most favoured regions [A,B,L]	786	86.6%
Residues in additional allowed regions [a,b,l,p]	112	12.3%
Residues in generously allowed regions [-a,-b,-l,-p]	7	0.8%
Residues in disallowed regions	3	0.3%


Number of non-glycine and non-proline residues	908	100.0%
Number of end-residues (excl. Gly and Pro)	12	
Number of glycine residues (shown as triangles)	92	
Number of proline residues	45	

Total number of residues	1057	

Based on an analysis of 118 structures of resolution of at least 2.0 Angstroms and R-factor no greater than 20%, a good quality model would be expected to have over 90% in the most favoured regions.

Ramachandran plots for all residue types


pdb1nek



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Ramachandran plots for all residue types


pdb1nek



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Ramachandran plots for all residue types


pdb1nek



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Chi1-Chi2 plots


pdb1nek



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0Å or better.

Chi1-Chi2 plots







pdb1nek



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Main-chain parameters

pdb1nek








Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. %-tage residues in A, B, L	908	86.6	74.8	10.0	1.2	BETTER
b. Omega angle st dev	1051	1.1	6.0	3.0	-1.6	BETTER
c. Bad contacts / 100 residues	83	7.9	12.2	10.0	-0.4	Inside
d. Zeta angle st dev	965	0.9	3.1	1.6	-1.4	BETTER
e. H-bond energy st dev	682	0.8	0.9	0.2	-0.7	Inside
f. Overall G-factor	1057	0.3	-0.6	0.3	3.0	BETTER

Side-chain parameters

pdb1nek




pdb1nek

Plot statistics


Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. Chi-1 gauche minus st dev	97	11.9	23.6	6.5	-1.8	BETTER
b. Chi-1 trans st dev	312	13.2	23.4	5.3	-1.9	BETTER
c. Chi-1 gauche plus st dev	420	13.4	22.0	4.9	-1.8	BETTER
d. Chi-1 pooled st dev	829	13.3	22.8	4.8	-1.9	BETTER
e. Chi-2 trans st dev	279	14.7	23.6	5.0	-1.8	BETTER

Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility



Key:- Helix Beta strand Random coil Accessibility shading: Buried Accessible


e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb1nek


a. Absolute deviation from mean Chi-1 value (excl. Pro)




b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors



Main-chain bond lengths

pdb1nek




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

Main-chain bond angles

pdb1nek




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

Main-chain bond angles

pdb1nek




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

RMS distances from planarity


pdb1nek



Histograms showing RMS distances of planar atoms from best-fit plane.
 Black bars indicate large deviations from planarity: RMS dist > 0.03 for rings, and > 0.02 otherwise.


Distorted geometry pdb1nek

Main-chain bond angles



Bond angles differing by > 10.0 degrees from small-molec values. Values shown: "ideal", actual, diff.

Planar groups



Sidechains with RMS dist. from planarity > 0.03Å for rings, or > 0.02Å otherwise. Value shown is RMS dist.