


# Ramachandran Plot

pdb1zp0




### Plot statistics

Residues in most favoured regions [A,B,L]	795	84.4%
Residues in additional allowed regions [a,b,l,p]	136	14.4%
Residues in generously allowed regions [-a,-b,-l,-p]	9	1.0%
Residues in disallowed regions	2	0.2%
-----		
Number of non-glycine and non-proline residues	942	100.0%
Number of end-residues (excl. Gly and Pro)	7	
Number of glycine residues (shown as triangles)	97	
Number of proline residues	46	
-----		
Total number of residues	1092	

Based on an analysis of 118 structures of resolution of at least 2.0 Angstroms and R-factor no greater than 20%, a good quality model would be expected to have over 90% in the most favoured regions.

# Ramachandran plots for all residue types


## pdb1zp0



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Ramachandran plots for all residue types


## pdb1zp0



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Ramachandran plots for all residue types


## pdb1zp0



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Chi1-Chi2 plots


## pdb1zp0



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Chi1-Chi2 plots







## pdb1zp0



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Main-chain parameters

## pdb1zp0








### Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. %-tage residues in A, B, L	942	84.4	53.6	10.0	3.1	BETTER
b. Omega angle st dev	1088	0.8	6.0	3.0	-1.7	BETTER
c. Bad contacts / 100 residues	28	2.6	32.6	10.0	-3.0	BETTER
d. Zeta angle st dev	995	0.8	3.1	1.6	-1.4	BETTER
e. H-bond energy st dev	696	0.8	1.2	0.2	-1.8	BETTER
f. Overall G-factor	1092	0.3	-1.0	0.3	4.4	BETTER

# Side-chain parameters

## pdb1zp0



pdb1zp0


### Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. Chi-1 gauche minus st dev	135	14.9	31.8	6.5	-2.6	BETTER
b. Chi-1 trans st dev	306	15.8	30.1	5.3	-2.7	BETTER
c. Chi-1 gauche plus st dev	419	14.3	28.8	4.9	-3.0	BETTER
d. Chi-1 pooled st dev	860	15.0	29.7	4.8	-3.0	BETTER
e. Chi-2 trans st dev	247	12.7	28.5	5.0	-3.2	BETTER




# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb1zp0


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb1zp0


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors






# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb1zp0


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb1zp0


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




Key:- Helix Beta strand Random coil Accessibility shading: Buried Accessible


e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors



# Main-chain bond lengths

## pdb1zp0




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

# Main-chain bond angles

## pdb1zp0




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

# Main-chain bond angles

## pdb1zp0




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

# RMS distances from planarity

## pdb1zp0




Histograms showing RMS distances of planar atoms from best-fit plane.  
 Black bars indicate large deviations from planarity: RMS dist > 0.03 for rings, and > 0.02 otherwise.



# Distorted geometry


## pdb1zp0

### Main-chain bond angles



Bond angles differing by > 10.0 degrees from small-molec values. Values shown: "ideal", actual, diff.

### Planar groups



Sidechains with RMS dist. from planarity > 0.03Å for rings, or > 0.02Å otherwise. Value shown is RMS dist.