


# Ramachandran Plot

pdb2acz




### Plot statistics

Residues in most favoured regions [A,B,L]	728	80.2%
Residues in additional allowed regions [a,b,l,p]	160	17.6%
Residues in generously allowed regions [-a,-b,-l,-p]	17	1.9%
Residues in disallowed regions	3	0.3%
Number of non-glycine and non-proline residues	908	100.0%
Number of end-residues (excl. Gly and Pro)	12	
Number of glycine residues (shown as triangles)	92	
Number of proline residues	45	
Total number of residues	1057	

Based on an analysis of 118 structures of resolution of at least 2.0 Angstroms and R-factor no greater than 20%, a good quality model would be expected to have over 90% in the most favoured regions.

# Ramachandran plots for all residue types


## pdb2acz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Ramachandran plots for all residue types


pdb2acz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Ramachandran plots for all residue types


## pdb2acz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Chi1-Chi2 plots


## pdb2acz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Chi1-Chi2 plots







## pdb2acz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Main-chain parameters

## pdb2acz








### Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. %-tage residues in A, B, L	908	80.2	64.1	10.0	1.6	BETTER
b. Omega angle st dev	1050	1.2	6.0	3.0	-1.6	BETTER
c. Bad contacts / 100 residues	55	5.2	22.3	10.0	-1.7	BETTER
d. Zeta angle st dev	964	1.0	3.1	1.6	-1.3	BETTER
e. H-bond energy st dev	675	0.8	1.1	0.2	-1.2	BETTER
f. Overall G-factor	1057	0.2	-0.8	0.3	3.4	BETTER

# Side-chain parameters

## pdb2acz



pdb2acz


### Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. Chi-1 gauche minus st dev	96	14.8	28.2	6.5	-2.1	BETTER
b. Chi-1 trans st dev	309	14.7	27.1	5.3	-2.3	BETTER
c. Chi-1 gauche plus st dev	423	15.4	25.8	4.9	-2.1	BETTER
d. Chi-1 pooled st dev	828	15.2	26.6	4.8	-2.3	BETTER
e. Chi-2 trans st dev	278	16.0	26.3	5.0	-2.1	BETTER




# Residue properties pdb2acz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2acz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2acz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2acz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2acz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2acz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2acz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2acz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors






# Residue properties pdb2acz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2acz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2acz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors



# Main-chain bond lengths

## pdb2acz




Black bars > 2.0 st. devs. from mean.

◀ or ▶ signifies data points off the graph in the direction shown.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

# Main-chain bond angles

## pdb2acz




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

# Main-chain bond angles

## pdb2acz




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

# RMS distances from planarity

## pdb2acz




Histograms showing RMS distances of planar atoms from best-fit plane.  
 Black bars indicate large deviations from planarity: RMS dist > 0.03 for rings, and > 0.02 otherwise.

# Distorted geometry


## pdb2acz

### Main-chain bond lengths




Bonds differing by > 0.05Å from small-molecule values. Values shown: "ideal", difference, actual

### Main-chain bond angles



Bond angles differing by > 10.0 degrees from small-molec values. Values shown: "ideal", actual, diff.

### Planar groups



Sidechains with RMS dist. from planarity > 0.03Å for rings, or > 0.02Å otherwise. Value shown is RMS dist.