


Ramachandran Plot

pdb2h89



Plot statistics

Residues in most favoured regions [A,B,L]	810	85.6%
Residues in additional allowed regions [a,b,l,p]	128	13.5%
Residues in generously allowed regions [-~a,-~b,-~l,-~p]	6	0.6%
Residues in disallowed regions	2	0.2%


Number of non-glycine and non-proline residues	946	100.0%
Number of end-residues (excl. Gly and Pro)	10	
Number of glycine residues (shown as triangles)	89	
Number of proline residues	50	

Total number of residues	1095	

Based on an analysis of 118 structures of resolution of at least 2.0 Angstroms and R-factor no greater than 20%, a good quality model would be expected to have over 90% in the most favoured regions.

Ramachandran plots for all residue types


pdb2h89



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Ramachandran plots for all residue types


pdb2h89



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Ramachandran plots for all residue types


pdb2h89



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Chi1-Chi2 plots


pdb2h89



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Chi1-Chi2 plots







pdb2h89



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

Main-chain parameters

pdb2h89




Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean
			Typical value	Band width	
a. %-tage residues in A, B, L	946	85.6	78.2	10.0	0.7 Inside
b. Omega angle st dev	1088	1.3	6.0	3.0	-1.6 BETTER
c. Bad contacts / 100 residues	97	8.9	9.0	10.0	0.0 Inside
d. Zeta angle st dev	1004	1.0	3.1	1.6	-1.3 BETTER
e. H-bond energy st dev	703	0.8	0.9	0.2	-0.7 Inside
f. Overall G-factor	1095	0.3	-0.6	0.3	2.8 BETTER

Side-chain parameters

pdb2h89




pdb2h89

Plot statistics


Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. Chi-1 gauche minus st dev	132	12.0	21.8	6.5	-1.5	BETTER
b. Chi-1 trans st dev	275	13.1	22.0	5.3	-1.7	BETTER
c. Chi-1 gauche plus st dev	451	12.5	20.5	4.9	-1.6	BETTER
d. Chi-1 pooled st dev	858	12.6	21.2	4.8	-1.8	BETTER
e. Chi-2 trans st dev	293	14.8	22.6	5.0	-1.6	BETTER

Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




c = cis-peptide

Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties pdb2h89


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors



Main-chain bond lengths

pdb2h89




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

Main-chain bond angles

pdb2h89




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

Main-chain bond angles

pdb2h89




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

RMS distances from planarity

pdb2h89




Histograms showing RMS distances of planar atoms from best-fit plane.
 Black bars indicate large deviations from planarity: RMS dist > 0.03 for rings, and > 0.02 otherwise.

▶ signifies data points off the graph in the direction shown.

Distorted geometry


pdb2h89

Main-chain bond angles



Bond angles differing by > 10.0 degrees from small-molec values. Values shown: "ideal", actual, diff.

Planar groups



Sidechains with RMS dist. from planarity > 0.03Å for rings, or > 0.02Å otherwise. Value shown is RMS dist.